

Oct. 23, 2009

Five-dimensional AGT Conjecture and the Deformed Virasoro Algebra

Hidetoshi Awata^{*} and Yasuhiko Yamada[†]

^{}Graduate School of Mathematics
Nagoya University, Nagoya 464-8602, Japan*

*[†]Department of Mathematics, Faculty of Science
Kobe University, Hyogo 657-8501, Japan*

Abstract

We study an analog of the AGT (Alday-Gaiotto-Tachikawa) relation in five dimensions. We conjecture that the instanton partition function of 5D $\mathcal{N} = 1$ pure $SU(2)$ gauge theory coincides with the inner product of the Gaiotto-like state in the deformed Virasoro algebra. In four-dimensional case, a relation between the Gaiotto construction and the theory of Braverman and Etingof is also discussed.

1 Introduction

In [1], Alday, Gaiotto and Tachikawa discovered remarkable relations between the 4D $\mathcal{N} = 2$ super conformal gauge theories and the 2D Liouville CFT. This surprising conjecture has been extended to asymptotic-free cases [2, 3] and to $SU(n)$ -gauge/ $\mathcal{W}_n$ -CFT [4, 5]. Some explanations were given from M -theory [6, 7] and from topological string [8]. Various checks have been done in [9, 10] for example. The AGT relation is also closely related with the current active subjects such as loop operators [11]–[13], wall crossing [14]–[16], integrability structures (see, e.g., [17, 18] and references therein) and so on.

In this paper, we will examine an extension of the AGT conjecture to the 5D gauge theory. The instanton counting [19]–[23] has a natural generalization to the 5D gauge theory [23, 24] that can be viewed as a q -analog of 4D cases. In [25]–[27] it was argued that they are closely related with the Macdonald symmetric polynomials [28]. On the other hand, there exists a natural deformation of the Virasoro/ $\mathcal{W}$ algebra [29]–[32] which is also related with Macdonald polynomials. Therefore, it is quite natural to expect an extension of the AGT relation between partition functions of the 5D gauge theory and correlation functions of the deformed Virasoro (or $\mathcal{W}$) algebra. In this note, we will formulate and check this relation in the simplest case: the pure $SU(2)$ gauge theory. Connections between 5D gauge theories and 2D relativistic integrable systems have been pointed out in [33, 34].

In section 2, following the work by Gaiotto [2], we will recall the conjectured relation between the instanton partition function of the 4D $\mathcal{N} = 2$ pure $SU(2)$ gauge theory and a Virasoro CFT correlation function (a two-point function or an inner product) with states of “irregular” singularity. In section 3, after a brief review on the deformed Virasoro algebra, we will formulate the 5D analog of the AGT conjecture in the simplest case: the pure $SU(2)$ gauge theory, and check it. We will see a very nontrivial coincidence also in 5D. Section 4 is devoted to conclusions and discussion. In the Appendix A, we will make a comment on the relation between the AGT conjecture and the instanton counting by Braverman and Etingof using the Whittaker-Toda equation [35, 36].

2 Review of Gaiotto construction

In the remarkable work [1], Alday, Gaiotto and Tachikawa conjectured the surprising relation between the 4D supersymmetric gauge theory and the 2D conformal field theory. In their original setting, it was conjectured that the instanton partition function Z^{inst} of

the 4D $\mathcal{N} = 2$ gauge theories with $N_f = 4$ flavours is coincides with the conformal block of the Virasoro CFT with four primary fields. The four fundamental matters corresponds to the four primary fields. Soon after that, Gaiotto gave extension of this correspondence to the theories with $N_f < 4$. These theories can be obtained as degeneration limit of the $N_f = 4$ case and certain confluent type operator/state appear. The state (which we call Gaiotto state) make irregular singularity for stress tensor $T(z)$ of order grater than two. Here, we will recall the construction of Gaiotto in case of $N_f = 0$.

Let V_h be the highest weight representation of the Virasoro algebra

$$[L_m, L_n] = (m - n)L_{m+n} + \frac{c}{12}(m^3 - m)\delta_{m+n,0}, \quad (2.1)$$

generated by the highest weight vector $|h\rangle$ such that

$$L_0|h\rangle = h|h\rangle, \quad L_n|h\rangle = 0 \quad (n > 0). \quad (2.2)$$

The dual representation V_h^* , and the pairing is defined by;

$$\langle h|L_0 = h\langle h|, \quad \langle h|L_n = 0 \quad (n < 0), \quad \langle h|h\rangle = 1. \quad (2.3)$$

We will put $c = 1 + 6(b + 1/b)^2$.

In order to realize the relation $\langle T(z) \rangle = \frac{\Lambda^2}{z^3} + \frac{2u}{z^2} + \frac{\Lambda^2}{z}$, which is the square of the Seiberg-Witten differential, Gaiotto defined a vector $|\Delta, \Lambda\rangle \in V_\Delta$ by the condition

$$L_1|\Delta, \Lambda\rangle = \Lambda^2|\Delta, \Lambda\rangle, \quad L_2|\Delta, \Lambda\rangle = 0, \quad \Lambda^2 \in \mathbb{C}, \quad (2.4)$$

with leading term normalization $|\Delta, \Lambda\rangle = |\Delta\rangle + \dots$ ¹ Note that (2.1) guarantees $L_n|\Delta, \Lambda\rangle = 0$ for all n larger than two.² Let $|\Delta, \Lambda\rangle = \sum_{n=0}^{\infty} \Lambda^{2n} v_n$ with $L_0 v_n = (\Delta + n)v_n$, $v_0 = |\Delta\rangle$, then (2.4) is equivalent to $L_1 v_n = v_{n-1}$ with $v_{-1} := 0$ and $L_2 v_n = 0$. The uniqueness of v_n is proved by the induction in n using the fact that the null vector $|\chi\rangle \in V_h$ defined by $L_n|\chi\rangle = 0$ for $n \geq 1$ does not exist for the generic h and c .

The dual vector $\langle \Delta, \Lambda| \in V_\Delta^*$ is defined similar way:

$$\langle \Delta, \Lambda|L_{-1} = \Lambda^2\langle \Delta, \Lambda|, \quad \langle \Delta, \Lambda|L_n = 0 \quad (n \leq -2), \quad (2.5)$$

$\langle \Delta, \Lambda| = \sum_{n=0}^{\infty} \Lambda^{2n} w_n$, where $w_n L_0 = (\Delta + n)w_n$ and $w_0 = \langle \Delta|$.

It is conjectured that the inner product $\langle \Delta, \Lambda|\Delta, \Lambda\rangle$ coincides with the instanton part of the 4D $\mathcal{N} = 2$ pure $SU(2)$ gauge theory. This conjecture has been checked for lower order in Λ -expansion and proved in the classical limit $c \rightarrow \infty$ [9]. In the appendix A, we will give another support on this conjecture by using the result of Braverman and Etingof [36].

¹Here $\dots$ means the higher order terms in L_0 -degree (level).

²Note also that if $|v\rangle \in V_h$ and $L_n|v\rangle = 0$ for $n = 2, 3, 4$ then $L_n L_1|v\rangle = L_n|v\rangle = 0$ for all $n \geq 2$.

3 Five-dimensional case

3.1 Quantum deformed Virasoro algebra

Let (q, t) be formal parameters. We also let $p := q/t$ and $t = q^\beta$. The q -Virasoro algebra is an associative algebra generated by $\langle T_n \rangle_{n \in \mathbb{Z}}$ with the relation [29, 30],

$$[T_n, T_m] = - \sum_{l=1}^{\infty} f_l (T_{n-l} T_{m+l} - T_{m-l} T_{n+l}) - \frac{(1-q)(1-t^{-1})}{1-p} (p^n - p^{-n}) \delta_{m+n,0}, \quad (3.6)$$

where $[A, B] = AB - BA$ and $f(z) = \sum_{l=0}^{\infty} f_l z^l$ with

$$f(x) = \exp \left\{ \sum_{n=1}^{\infty} \frac{(1-q^n)(1-t^{-n})}{1+p^n} \frac{x^n}{n} \right\}. \quad (3.7)$$

The defining relation (3.6) can be written in terms of the current $T(z) = \sum_{n \in \mathbb{Z}} T_n z^{-n}$ as

$$f(w/z) T(z) T(w) - T(w) T(z) f(z/w) = - \frac{(1-q)(1-t^{-1})}{1-p} \left\{ \delta \left(\frac{pw}{z} \right) - \delta \left(\frac{w}{pz} \right) \right\}, \quad (3.8)$$

where $\delta(z) = \sum_{n \in \mathbb{Z}} z^n$ is the multiplicative δ -function which satisfies $g(z) \delta(z) = g(1) \delta(z)$ for any function $g(z) = \sum_{n \in \mathbb{Z}} g_n z^n$.

The q -Virasoro algebra reduces to the Virasoro algebra in the limit $q \rightarrow 1$. Let $q = e^{\hbar}$. Then the generator $L(z)$ defined as

$$T(z) = 2 + \beta \hbar^2 \left\{ z^2 L(z) + \frac{1}{4} \left(\sqrt{\beta} - 1/\sqrt{\beta} \right)^2 \right\} + O(\hbar^4), \quad (3.9)$$

is the ordinary Virasoro current $L(z) = \sum_{n \in \mathbb{Z}} L_n z^{-n-2}$, which satisfies (2.1) with the central charge $c = 1 - 6 \left(\sqrt{\beta} - 1/\sqrt{\beta} \right)^2$, $(\beta = -b^2)$.

3.2 Deformed Gaiotto state

Let M_h be the Verma module over the q -Virasoro algebra, generated by the highest weight vector $|h\rangle$, such that $T_n |h\rangle = 0$ for $n > 0$ and $T_0 |h\rangle = h |h\rangle$ with $h \in \mathbb{C}$. The dual module M_h^* is generated by $\langle h|$ which satisfies $\langle h| T_n = 0$ for $n < 0$ and $\langle h| T_0 = h \langle h|$. The bilinear form $M_h^* \otimes M_h \rightarrow \mathbb{C}$ is uniquely defined by $\langle h|h\rangle = 1$. For a generic value of q , the representation theory of the q -Virasoro algebra is similar to that of the Virasoro algebra [37]. Let us introduce the (outer) grading operator d which satisfies $[d, T_n] = n T_n$ and set $d|h\rangle = 0$. We call a vector $|v\rangle \in M(h)$ of level n if $d|v\rangle = -n|v\rangle$. Then M_h has the grade decomposition $M_h = \oplus_{n=0}^{\infty} M_{h,n}$ by the grade n subspace $M_{h,n}$.

The deformed analog of the Gaiotto state $|G\rangle \in M_h$ is defined by

$$T_1|G\rangle = \Lambda^2|G\rangle, \quad T_n|G\rangle = 0 \quad (n \geq 2), \quad \Lambda^2 \in \mathbb{C}, \quad (3.10)$$

with a normalization condition $|G\rangle = |h\rangle + \dots$. Let $|G\rangle = \sum_{n=0}^{\infty} \Lambda^{2n}|G_n\rangle$ with $|G_n\rangle \in M_{h,n}$, then (3.10) is equivalent to $T_1|G_n\rangle = |G_{n-1}\rangle$ with $|G_{-1}\rangle := 0$ and $T_m|G_n\rangle = 0$ for $m \geq 2$. The dual vector $\langle G| \in M_h^*$ is defined similar way:

$$\langle G|T_{-1} = \Lambda^2\langle G|, \quad \langle G|T_n = 0 \quad (n \leq -2), \quad (3.11)$$

and $\langle G| = \langle h| + \dots$. To calculate an inner product $\langle G|G\rangle = \sum_{n=0}^{\infty} \Lambda^{4n}\langle G_n|G_n\rangle$, it is convenient to introduce a free boson realization for the q -Virasoro algebra.

3.3 Free boson realization

The q -Virasoro algebra has a free field representation:

$$T(z) = \sum_{n \in \mathbb{Z}} T_n z^{-n} = \Lambda_1(z) + \Lambda_2(z), \quad (3.12)$$

$$\Lambda_1(z) = \exp \left\{ - \sum_{n=1}^{\infty} t^n a_{-n} z^n \right\} \exp \left\{ \sum_{n=1}^{\infty} q^n a_n z^{-n} \right\} K, \quad (3.13)$$

$$\Lambda_2(z) = \exp \left\{ \sum_{n=1}^{\infty} q^n a_{-n} z^n \right\} \exp \left\{ - \sum_{n=1}^{\infty} t^n a_n z^{-n} \right\} K^{-1}. \quad (3.14)$$

Where a_n are the Heisenberg operators realized as

$$a_n = \frac{q^n - 1}{t^n} \frac{\partial}{\partial p_n}, \quad a_{-n} = v_n p_n, \quad v_n := \frac{t^n - 1}{n(t^n + q^n)q^n} \quad (n > 0), \quad (3.15)$$

with the variables $p = (p_1, p_2, \dots)$ and K is a multiplicative charge operator which commute with a_n 's. Let $|k\rangle$ be the highest weight vector such as $K|k\rangle = k|k\rangle$ and $a_n|k\rangle = 0$ ($n > 0$). Then $T_0|k\rangle = (k + k^{-1})|k\rangle$. The deformed Gaiotto vector is defined by (3.10) and $|G\rangle = |k\rangle + \dots$. This over determined system of linear equations have unique solution.³ The vector $|G\rangle$ is realized as a polynomial in $p = (p_1, p_2, \dots)$

$$G(p) = 1 + \Lambda^2 c_1 p_1 + \Lambda^4 (c_{11} p_1^2 + c_2 p_2) + \dots, \quad (3.16)$$

³If v_n 's are considered as free parameters without the relations (3.15), the equations become inconsistent at level 5. It may be interesting to study the general form of the parameters v_n which admits the Gaiotto like state.

The first few coefficients are as follows;

$$\begin{aligned}
c_1 &= \frac{kt}{(k^2q - t)(q - 1)}, \\
c_{11} &= \frac{k^2(kq - t)(kq + t)t^2}{2(k^2q^2 - t)(k^2q - t)(k^2q - t^2)(q - 1)^2}, \\
c_2 &= -\frac{k^2(k^2q^2 + t^2)t^2}{2(k^2q^2 - t)(k^2q - t)(k^2q - t^2)(q^2 - 1)}, \\
c_{111} &= \frac{k^3(k^4q^5 - 2k^2t^3q^3 + k^2t^2q^3 + k^2t^3q^2 - 2k^2t^2q^2 + t^5)t^3}{6(k^2q^3 - t)(k^2q^2 - t)(k^2q - t)(k^2q - t^2)(k^2q - t^3)(q - 1)^3}, \\
c_{21} &= -\frac{k^3(k^2q^3 - t^3)(k^2q^2 + t^2)t^3}{2(k^2q^3 - t)(k^2q^2 - t)(k^2q - t)(k^2q - t^2)(k^2q - t^3)(q - 1)(q^2 - 1)}, \\
c_3 &= \frac{k^3(k^4q^5 + k^2t^3q^3 + k^2t^2q^3 + k^2t^3q^2 + k^2t^2q^2 + t^5)t^3}{3(k^2q^3 - t)(k^2q^2 - t)(k^2q - t)(k^2q - t^2)(k^2q - t^3)(q^3 - 1)}. \tag{3.17}
\end{aligned}$$

As in [29], one can identify the boson Fock space with the space of symmetric functions in variables, say $x = (x_1, x_2, \dots)$, by the natural identification $p_n = \sum_i x_i^n$. Then the function $G(p)$ is written as a sum of the Macdonald function $P_\lambda(x; q, t)$ [28] as

$$G(p) = \sum_{\lambda} \Lambda^{2|\lambda|} P_\lambda(x; q, t) \prod_{(i,j) \in \lambda} \frac{k}{1 - k^2 q^j t^{-i}} \frac{q^{\lambda_i - j}}{1 - q^{\lambda_i - j + 1} t^{\lambda_j^\vee - i}}, \tag{3.18}$$

which we have checked for $|\lambda| \leq 8$. Similar structure can be observed in 4D case in terms of the Jack polynomials.

Next, we compute the dual vector $\langle G|$ defined by (3.11). To do this, we use an involution ι such that

$$\iota(AB) = \iota(B)\iota(A), \quad \iota(T_n) = T_{-n}, \quad \iota(a_n) = a_{-n}, \quad \iota(K) = K^{-1}. \tag{3.19}$$

Then, the free field realization of the vector $\langle G|$ is obtained from the polynomial $G(p)$ simply by the replacement

$$p_n \rightarrow u_n \frac{\partial}{\partial p_n}, \quad u_n := -n(t^n + q^n) \left(\frac{q}{t}\right)^n \frac{1 - q^n}{1 - t^n}, \quad k \rightarrow k^{-1}. \tag{3.20}$$

Hence the inner product is given as

$$\langle G|G \rangle = \sum_{\lambda} \Lambda^{4|\lambda|} c_{\lambda}(k) c_{\lambda}(k^{-1}) w_{\lambda}, \tag{3.21}$$

where for a partition $\lambda = (\lambda_1, \lambda_2, \dots, \lambda_{\ell}) = (1^{m_1} 2^{m_2} \dots |\lambda|^{m_{|\lambda|}})$,

$$w_{\lambda} := \prod_{i=1}^{|\lambda|} m_i! u_i^{m_i}. \tag{3.22}$$

Then we have

$$\langle G|G \rangle = 1 + \Lambda^4 k^2 \frac{qt(q+t)}{(q-1)(k^2q-t)(q-tk^2)(1-t)} + \Lambda^8 k^4 (qt)^2 \frac{A}{B} + \dots, \quad (3.23)$$

where

$$\begin{aligned} A = & k^2 q^7 + k^2 t^2 q^6 + 2k^2 t q^6 - k^4 t^3 q^5 - t^3 q^5 + 2k^2 t^2 q^5 + k^2 t q^5 \\ & - k^4 t^4 q^4 - t^4 q^4 - k^4 t^3 q^4 + k^2 t^3 q^4 - t^3 q^4 - k^4 t^2 q^4 - t^2 q^4 \\ & - k^4 t^5 q^3 - t^5 q^3 - k^4 t^4 q^3 + k^2 t^4 q^3 - t^4 q^3 - k^4 t^3 q^3 - t^3 q^3 \\ & + k^2 t^6 q^2 + 2k^2 t^5 q^2 - k^4 t^4 q^2 - t^4 q^2 + 2k^2 t^6 q + k^2 t^5 q + k^2 t^7, \\ B = & (q^2 - 1)(q - 1)(k^2 q^2 - t)(k^2 q - t)(k^2 q - t^2)(1 - t)(1 - t^2) \\ & \times (q^2 - tk^2)(q - tk^2)(q - t^2 k^2). \end{aligned} \quad (3.24)$$

This is the 5D analog of $\langle \Delta, \Lambda | \Delta, \Lambda \rangle$ which can be compared with the 5D instanton partition function.

The instanton part of the Nekrasov's five-dimensional pure $SU(2)$ partition function is written by a sum over two Young diagrams λ and μ as follows [24];⁴

$$\begin{aligned} Z^{\text{inst}} = & \sum_{\lambda, \mu} \frac{(\Lambda^4 t / q)^{|\lambda| + |\mu|}}{N_{\lambda\lambda}(1) N_{\lambda\mu}(Q) N_{\mu\mu}(1) N_{\mu\lambda}(Q^{-1})}, \\ N_{\lambda\mu}(Q) := & \prod_{(i,j) \in \mu} \left(1 - Q q^{\lambda_i - j} t^{\mu_j^\vee - i + 1} \right) \prod_{(i,j) \in \lambda} \left(1 - Q q^{-\mu_i + j - 1} t^{-\lambda_j^\vee + i} \right). \end{aligned} \quad (3.25)$$

Here, $\lambda = (\lambda_1, \lambda_2, \dots)$ is a Young diagram/partition such that $\lambda_i \geq \lambda_{i+1}$. $\lambda^\vee$ is its conjugate (dual) as Young diagram and $|\lambda| = \sum_i \lambda_i$. Then we observe the following coincidence:

$$Z^{\text{inst}} = \langle G|G \rangle, \quad (3.26)$$

for $k = Q^{\frac{1}{2}}$. We have checked this up to level 9.

4 Summary and discussion

We studied 5D analog of AGT conjecture in the simplest pure $SU(2)$ gauge theory. It turns out that a natural generalization of the Gaiotto construction using the deformed Virasoro algebra is related to the gauge theory. Fortunately, the parameters (q, t) are directly related in both sides of the correspondence. This is because these parameters

⁴The parameter Q is related with the vacuum expectation value a of the scalar fields in the vector multiplets as $Q = q^a$ and the parameters (q, t) are related with those (ϵ_1, ϵ_2) of the Ω background through $(q, t) = (e^{\epsilon_2}, e^{-\epsilon_1})$.

have already been chosen in order to fit the conventions in Macdonald polynomials. It is quite natural to expect similar relation between $SU(n)$ gauge theory and the deformed $\mathcal{W}_n$ algebras [31, 32]. Inclusion of matter fields is also interesting problem. For the proof of the conjecture, the vertex operator constructions [23, 38, 39] would be useful.

In 4D case, the Gaiotto states produce irregular singularities. Such a conformal field theory with irregular singularities [40] are recently studied from the point of view of quantum Painlevé equations. From this point of view, the AGT relation may be considered as a quantization of the correspondence between the Seiberg-Witten theory and the Hitchin/Painlevé equations (see [41] for example).

Acknowledgments

We would like thank H. Kanno, H. Nakajima and Y. Tachikawa for discussions. The work of Y.Y. is supported in part by Grant-in-Aid for Scientific Research [#21340036] from the Japan Ministry of Education, Culture, Sports, Science and Technology. The work of H.A. is supported in part by Daiko Foundation.

Appendix A: Relation to the Braverman and Etingof

We follow the notation in section 2. Let us consider the one-point function

$$\Psi(z) = \langle \Delta_2, \Lambda | \Phi_h(z) | \Delta_1, \Lambda \rangle. \quad (\text{A.1})$$

We will derive the 2nd order differential equation for $\Psi(z)$ when the operator $\Phi_h(z)$ is the degenerate primary field of conformal dimension $h = h_{2,1}$. To do this, we consider the insertion of the stress tensor $T(w)$:

$$\langle \Delta_2, \Lambda | T(w) \Phi_h(z) | \Delta_1, \Lambda \rangle = \sum_{n \in \mathbf{Z}} \langle \Delta_2, \Lambda | L_n w^{-n-2} \Phi_h(z) | \Delta_1, \Lambda \rangle. \quad (\text{A.2})$$

Using the defining relations of the states $\langle \Delta_2, \Lambda |$, $|\Delta_1, \Lambda\rangle$ and the operator $\Phi_h(z)$:

$$[L_n, \Phi_h(z)] = z^n \left[z \frac{d}{dz} + h(n+1) \right] \Phi_h(z), \quad (\text{A.3})$$

one can derive the following relation

$$\langle \Delta_2, \Lambda | T(w) \Phi_h(z) | \Delta_1, \Lambda \rangle = \left[\frac{\Lambda^2}{w} + \frac{\Lambda^2}{w^3} + \left(\frac{1}{w-z} - \frac{1}{w} \right) \frac{\partial}{\partial z} + \frac{h}{(w-z)^2} \right] \Psi(z)$$

$$+\frac{1}{w^2}\langle\Delta_2, \Lambda|\Phi_h(z)L_0|\Delta_1, \Lambda\rangle. \quad (\text{A.4})$$

On the other hand, by noting the relation

$$\frac{1}{2}\Lambda\frac{\partial}{\partial\Lambda}|\Delta_1, \Lambda\rangle = (L_0 - \Delta_1)|\Delta_1, \Lambda\rangle, \quad \frac{1}{2}\Lambda\frac{\partial}{\partial\Lambda}\langle\Delta_2, \Lambda| = \langle\Delta_2, \Lambda|(L_0 - \Delta_2), \quad (\text{A.5})$$

we have

$$\langle\Delta_2, \Lambda|\Phi_h(z)L_0|\Delta_1, \Lambda\rangle = \left[\frac{\Lambda}{4}\frac{\partial}{\partial\Lambda} + \frac{\Delta_1 + \Delta_2 - h}{2} - \frac{z}{2}\frac{\partial}{\partial z}\right]\Psi(z). \quad (\text{A.6})$$

Thus we get the following formula

$$\begin{aligned} \langle\Delta_2, \Lambda|T(w)\Phi_h(z)|\Delta_1, \Lambda\rangle &= \left[\frac{\Lambda^2}{w} + \frac{\Lambda^2}{w^3} + \left(\frac{1}{w-z} - \frac{1}{w}\right)\frac{\partial}{\partial z} + \frac{h}{(w-z)^2}\right. \\ &\quad \left.+ \frac{1}{w^2}\left(\frac{\Lambda}{4}\frac{\partial}{\partial\Lambda} + \frac{\Delta_1 + \Delta_2 - h}{2} - \frac{z}{2}\frac{\partial}{\partial z}\right)\right]\Psi(z). \end{aligned} \quad (\text{A.7})$$

This gives the correlation function of the 2nd descendant operator $\widehat{L}_{-2}\Phi_h(z)$ defined by

$$T(w)\Phi_h(z) = \sum_{n \in \mathbf{Z}} (w-z)^{-n-2} \widehat{L}_n \Phi_h(z), \quad (\text{A.8})$$

as follows

$$\langle\Delta_2, \Lambda|\widehat{L}_{-2}\Phi_h(z)|\Delta_1, \Lambda\rangle = \frac{1}{z^2}\left[\Lambda^2\left(z + \frac{1}{z}\right) - \frac{3}{2}z\frac{\partial}{\partial z} + \frac{\Lambda}{4}\frac{\partial}{\partial\Lambda} + \frac{\Delta_1 + \Delta_2 - h}{2}\right]\Psi(z). \quad (\text{A.9})$$

When the primary field Φ_h has the dimension

$$h = h_{2,1} = -\frac{1}{2} - \frac{3}{4b^2}, \quad (\text{A.10})$$

and the null vector at level 2

$$(b^2\widehat{L}_{-1}^2 + \widehat{L}_{-2})\Phi_h(z) = 0, \quad (\text{A.11})$$

we obtain the following differential equation

$$\left[b^2z^2\frac{\partial^2}{\partial z^2} + \Lambda^2\left(z + \frac{1}{z}\right) - \frac{3}{2}z\frac{\partial}{\partial z} + \frac{\Lambda}{4}\frac{\partial}{\partial\Lambda} + \frac{\Delta_1 + \Delta_2 - h}{2}\right]\Psi(z) = 0. \quad (\text{A.12})$$

According to the fusion rule $[h_{2,1}] \otimes [\Delta(a)] = [\Delta(a \pm \frac{1}{2b})]$ where $\Delta(a) = (b+1/b)^2/4 - a^2$, we will take $\Delta_1 = \Delta(a + \frac{1}{4b})$ and $\Delta_2 = \Delta(a - \frac{1}{4b})$. Then, by putting $\Psi(z) = z^{\Delta_2 - \Delta_1 - h_{2,1}}Y(x)$ and $z = e^x$, we finally obtain the following differential equation for $Y(x)$

$$\left[b^2\frac{\partial^2}{\partial x^2} + 2ab\frac{\partial}{\partial x} + \Lambda^2(e^x + e^{-x}) + \frac{\Lambda}{4}\frac{\partial}{\partial\Lambda}\right]Y(x) = 0. \quad (\text{A.13})$$

This equation coincides with the differential equation of Braverman-Etingof (eq(2.10) [36]) whose WKB like solution⁵ gives the instanton part of the 4D $\mathcal{N} = 2$ pure $SU(2)$ gauge theory with equivalent parameters $(\epsilon_1, \epsilon_2) = (b, 0)$. Note that in the AGT conjecture, one put $(\epsilon_1, \epsilon_2) = (b, \frac{1}{b})$. The difference corresponds to the charge shift in $\langle \Delta(a - \frac{1}{4b}), \Lambda | \Phi_h(z) | \Delta(a + \frac{1}{4b}), \Lambda \rangle$. It will be interesting to study the 5D difference analog of these differential equations, which may be related with the relativistic Toda system [33, 34].

References

- [1] L. Alday, D. Gaiotto and Y. Tachikawa, “Liouville Correlation Functions from Four-dimensional Gauge Theories,” [arXiv:0906.3219](#).
- [2] D. Gaiotto, “Asymptotically free $N = 2$ theories and irregular conformal blocks,” [arXiv:0908.0307](#).
- [3] A. Marshakov, A. Mironov and A. Morozov, *Phys. Lett.* **B682** (2009) 125–129, [arXiv:0909.2052](#).
- [4] N. Wyllard, *JHEP* **0911** (2009) 002, [arXiv:0907.2189](#).
- [5] A. Mironov and A. Morozov, *Nucl. Phys.* **B825** (2009) 1–37, [arXiv:0908.2569](#).
- [6] G. Bonelli and A. Tanzini, “Hitchin systems, $N = 2$ gauge theories and W-gravity,” [arXiv:0909.4031](#).
- [7] L. Alday, F. Benini and Y. Tachikawa, “Liouville/Toda central charges from M5-branes,” [arXiv:0909.4776](#).
- [8] R. Dijkgraaf and C. Vafa, “Toda Theories, Matrix Models, Topological Strings, and $N = 2$ Gauge Systems,” [arXiv:0909.2453](#).
- [9] A. Mironov and A. Morozov, *Phys. Lett.* **B682** (2009) 118–124, [arXiv:0909.3531](#).
- [10] R. Poghossian, “Recursion relations in CFT and $N = 2$ SYM theory,” [arXiv:0909.3412](#).
- [11] N. Drukker, D. Morrison and T. Okuda, *JHEP* **0909** (2009) 031, [arXiv:0907.2593](#).

⁵The WKB parameter κ in front of $\Lambda \frac{\partial}{\partial \Lambda}$ is hidden in our normalization.

- [12] L. Alday, D. Gaiotto, S. Gukov, Y. Tachikawa and H. Verlinde, “Loop and surface operators in $N = 2$ gauge theory and Liouville modular geometry,” [arXiv:0909.0945](#).
- [13] N. Drukker, J. Gomis, T. Okuda and J. Teschner, “Gauge Theory Loop Operators and Liouville Theory,” [arXiv:0909.1105](#).
- [14] D. Gaiotto, G. Moore, A. Neitzke, “Wall-crossing, Hitchin Systems, and the WKB Approximation,” [arXiv:0907.3987](#).
- [15] S. Cecotti and C. Vafa, “BPS Wall Crossing and Topological Strings,” [arXiv:0910.2615](#).
- [16] M. Kontsevich and Y. Soibelman, “Stability structures, motivic Donaldson-Thomas invariants and cluster transformations,” [arXiv:0811.2435](#).
- [17] K. Takasaki, *J. Geom. Phys.* **59** (2009) 1244–1257, [arXiv:0903.2607](#).
- [18] N. Nekrasov and S. Shatashvili, “Quantization of Integrable Systems and Four Dimensional Gauge Theories,” [arXiv:0908.4052](#).
- [19] N. Nekrasov, *Adv. Theor. Math. Phys.* **7** (2003) 831–864, [arXiv:hep-th/0206161](#).
- [20] R. Flume and R. Poghossian, *Int. J. Mod. Phys.* **A18** (2003) 2541, [arXiv:hep-th/0208176](#).
- [21] U. Bruzzo, F. Fucito, J. Morales and A. Tanzini, *JHEP* **0305** (2003) 054, [arXiv:hep-th/0211108](#).
- [22] H. Nakajima and K. Yoshioka, *Invent. Math.* **162** (2005) 313–355, [arXiv:math.AG/0306198](#).
- [23] N. Nekrasov and A. Okounkov, *The unity of mathematics*, Progr. Math. **244**, Birkhäuser, Boston, 2006, 525–596, [arXiv:hep-th/0306238](#).
- [24] H. Nakajima and K. Yoshioka, *Transform. Groups* **10** (2005), 489–519, [arXiv:math.AG/0505553](#).
- [25] H. Awata and H. Kanno, *JHEP* **0505** (2005) 039, [arXiv:hep-th/0502061](#).
- [26] A. Iqbal, C. Kozcaz and C. Vafa, *JHEP* **0910** (2009) 069, [arXiv:hep-th/0701156](#).
- [27] H. Awata and H. Kanno, *Int. J. Mod. Phys.* **A24** (2009) 2253–2306, [arXiv:0805.0191](#).

- [28] I.G. Macdonald, *Symmetric functions and Hall polynomials*, Second Edition, Oxford University Press, 1995.
- [29] J. Shiraishi, H. Kubo, H. Awata and S. Odake, *Lett. Math. Phys.* **38**, 33–51, [arXiv:q-alg/9507034](#).
- [30] H. Awata, H. Kubo, S. Odake and J. Shiraishi, Les Publications CRM, *Proceedings of the Canada-China meeting on theoretical physics*, (Edited by L. Lapointe et.al.) 2003, pp. 154–188, [arXiv:hep-th/9612233](#).
- [31] B. Feigin and E. Frenkel, *Commun. Math. Phys.* **178** (1996) 653–678, [arXiv:q-alg/9508009](#).
- [32] H. Awata, H. Kubo, S. Odake and J. Shiraishi, *Commun. Math. Phys.* **179** (1996) 401–416, [arXiv:q-alg/9508011](#).
- [33] A. Gorsky and N. Nekrasov, *Nucl. Phys.* **B436** (1995) 582–608, [arXiv:hep-th/9401017](#).
- [34] N. Nekrasov, *Nucl. Phys.* **B531** (1998) 323–344, [arXiv:hep-th/9609219](#).
- [35] A. Braverman, “Instanton counting via affine Lie algebras I: Equivariant J-functions of (affine) flag manifolds and Whittaker vectors,” [arXiv:math/0401409](#).
- [36] A. Braverman and P. Etingof, “Instanton counting via affine Lie algebras II: from Whittaker vectors to the Seiberg-Witten prepotential,” [arXiv:math/0409441](#).
- [37] P. Bouwknegt and K. Pilch, *Commun. Math. Phys.* **196** (1998) 249–288, [arXiv:q-alg/9710026](#).
- [38] T. Maeda, T. Nakatsu, K. Takasaki and T. Tamakoshi, *JHEP* **0503** (2005) 056, [arXiv:hep-th/0412327](#).
- [39] E. Carlsson, “Vertex Operators and Moduli Space of Sheaves,” [arXiv:math/0906.1825](#).
- [40] M. Jimbo, H. Nagoya and J. Sun, *J. Phys. A: Math. Theor.* **41** (2008) 175205 (14pp).
- [41] K. Kajiwara, T. Masuda, M. Noumi, Y. Ohta and Y. Yamada, *Funkcial. Ekvac.* **48** (2005) 147–160, [arXiv:nlin/0403009](#).